

## ***Maro lehtineni* (Araneae: Linyphiidae) – a spider species new to the fauna of Poland**

**Konrad Wiśniewski & Wanda Wesołowska**

doi: 10.5431/aramit5011

**Abstract.** The rare linyphiid spider species *Maro lehtineni* Saaristo, 1971 was recorded in the Polish part of the Sudetes, in the Giant Mountains (in Polish Karkonosze). Five males were found on one of the sloping transition mires in the spring and early summer of 2011 and 2012. We provide new figures for identification of this species, and summarize and discuss data on its distribution, characteristic habitats and phenology.

**Keywords:** Giant Mountains, mires, mountains, Sudetes

**Zusammenfassung.** *Maro lehtineni* (Araneae: Linyphiidae) – eine neue Spinnenart für die Fauna von Polen. Die seltene Linyphiidenart *Maro lehtineni* Saaristo, 1971 konnte im polnischen Teil des Riesengebirges (polnisch Karkonosze) nachgewiesen werden. Fünf Männchen wurden auf einem Hangzwischenmoor gefunden. Wir veröffentlichen neue Abbildungen zur Bestimmung dieser Art und fassen die Daten über Verbreitung, bevorzugten Lebensraum und Phänologie zusammen.

*Maro* O. P.-Cambridge, 1906 is a genus of very small spiders belonging to Linyphiidae which includes 17 species (World Spider Catalog 2015) living in the Palaearctic and the northern Nearctic. There are six European species (Tanasevitch & Nekhaeva 2014, van Helsdingen 2015); four species have been found so far in central Europe. All *Maro* spiders, apart from being minute, seem to have secluded way of life and are therefore rarely found. However, it was recently shown that at least two of them – *Maro lepidus* Casimir, 1961 and *Maro minutus* O. P.-Cambridge, 1906 – might be quite numerous on mountainous mires (Wiśniewski & Wesołowska 2012).

Our knowledge of the distribution of *Maro* in Poland has increased greatly in the last three decades. Only *M. minutus* is widely distributed (Kupryjanowicz 2008) and has been recently recorded from different localities (Sudetes – Woźny & al. 1988, Masurian Lakeland – Staręga 1996, Polesie – Staręga 2000, Poleski National Park – Stańska et al. 2002, Biebrzański National Park – Kupryjanowicz 2003). *Maro sublestus* Falconer, 1915 was found in the Sudetes, in the Giant Mountains (Karkonosze/ Riesengebirge) by Szymkowiak (2004). Wiśniewski & Wesołowska (2012) documented the presence of

*M. lepidus* in different ranges of the Sudetes (Giant Mountains, Iżera Mountains, Table Mountains), but also confirmed and supplemented data on distribution with regards to *M. minutus* and *M. sublestus* (Giant Mountains, Iżera Mountains).

The record of another species from this genus – *Maro lehtineni* – is presented here, thus all four species known from central Europe to date are also present in Poland, in the Giant Mountains. Our main aims are to provide new figures, which might be helpful for species identification, and summarize data on the biology of *M. lehtineni*.

### **Material and methods**

The material comes from a study in which the diversity of spider assemblages in mountainous mires in the Sudetes was analysed. Spiders were sampled using pitfall traps and sweep nets in 21 mountainous mires in the Polish part of the Giant Mountains (Karkonosze) and in the Iżera Mountains from April to October in 2010 and 2011, and from April to May in 2012. Mountainous raised bogs and poor fens were investigated. The other study that concerned selected habitats surrounding four of these mires was carried out in 2012; however no specimens of *M. lehtineni* were recorded there.

Five males of *M. lehtineni* were found in a single transition mire, on the slopes of the Giant Mountains (50°46.075'N, 15°41.255'E; 1210m a.s.l.; Figs 1, 2). All *M. lehtineni* were caught using pitfall traps, in the spring and early summer (trap exposure peri-

Konrad WIŚNIEWSKI & Wanda WESOŁOWSKA, University of Wrocław, Department of Biodiversity and Evolutionary Taxonomy, Przybyszewskiego 63/77, 51-148 Wrocław; E-mail: konwisniew@gmail.com, tomwes@biol.uni.wroc.pl



**Figs 1-2:** Mire where *Maro lehtineni* was recorded

od for these records: 27 April–12 May 2011, 12–30 May 2011, 16 June–3 July 2011, 10–29 May 2012). This species cohabited with *M. lepidus* (21 males, 4 females) and *M. sublestus* (1 male) in the same mire.


### Identification of *Maro lehtineni*

The three species already known from Poland (*M. lepidus*, *M. minutus* and *M. sublestus*) are fairly easy to identify based on their lamella characteristic in the male palp and general appearance of the female epigyne (Saaristo 1971). However, recognising *M. lehtineni* requires more care.

Males of *M. lehtineni* (Fig. 3) are distinguished from congeners by the shape of the lamella charac-

teristica, which is not divided (as in *M. minutus*), but pointed with hardly visible denticles (compared to *M. sublestus* which has this part of the bulb larger, truncated and with a conspicuously indentated apical margin), thin and not scale-like (distinguishing it from *M. lepidus*). The overall appearance of this structure is thus characteristic, but due to its extremely small size it may be difficult to see the differences immediately. Furthermore, its shape is strongly dependent on the angle, from which the palp is being observed and possibly fairly changeable (Figs 4–8, compare also Saaristo 1971, Thaler 1983, Bolzern et al. 2005). In telling apart the four species, it is also useful to compare the overall proportions of the paracymbium.





**Figs 3-5:** Male of *Maro lehtineni*; **3.** general appearance, **4.-5.** palp in lateral view, seen from slightly different angles.

Female genitalia of *M. lehtineni* are similar to those of *M. lepidus*. Unfortunately during the study we did not find any female of the newly recorded species. The figures and descriptions by Saaristo (1971) offer a sufficient basis for distinguishing females of the two species.

### Distribution, habitats and phenology

*Maro lehtineni* has not been recorded in Poland before, but the single specimen of this species was found fairly close to the Polish border, in the Czech part of the Giant Mountains (Růžička 2000). Apart from the Czech Republic it is also known from Austria (Thaler 1983), Belgium (Baert & Kekenbosch 1979), Germany (Blick et al. 2004), Finland (Saaristo 1971), Norway (Hauge 1980), Slovakia (Svatoň & Gajdoš 2004), Sweden (Granström 1978) and Switzerland (Bolzern et al. 2005).


There is little information about typical habitats and time of occurrence of the presented species, because data are very scarce. In this study all specimens were found in a typical, sloping, transition mire in the upper-forest zone of the Giant Mountains (where it ranges app. from 1000 to 1250 m a.s.l.). This habitat seems to be characteristic for *M. lepidus* instead, which is sometimes regarded as tyrphobiont species (Casemir 1976) and indeed this species was the most numerous *Maro* in this mire. In addition, females – which are less active than males – of *M. lepidus* were recorded there. The presence of *M. lehtineni* might also be connected with its dispersal from the other habitats (in case of this particular

mire surrounding habitats were not analysed) and it is hard to justify here what its real affinity towards mountainous mires is.

Previous records of this species came from the northern Europe, where it has been recorded among moss *Hylocomium* (Saaristo 1971), in *Sphagnum* (Palmgren 1975), both within spruce forests, or in *Calluna* heaths with *Hylocomium* (Hauge 1980). Moreover, the species was found in different habitats in the mountains, such as hay meadows (Thaler 1983), dwarf-shrub heaths (Thaler 1983, Bolzern et al. 2005, Muff et al. 2007), timberline and spruce forest (Muff et al. 2007). It was also found in the other places with suitable habitats, including bogs or fens (Baert & Kekenbosch 1979, Svatoň & Gajdoš 2004, Martin 2009), sedge swamp or swamp forests (Martin 2009). In summary, it was found in the variety of habitats in northern Europe and in the mountains, but also in some places with high humidity in the lowlands of Central Europe.

*M. lehtineni* was found predominately in early spring, just after the snow melted, however one male was observed later, in early summer. This activity pattern resembles that of *M. lepidus* (see Wiśniewski & Wesołowska 2012). In the other studies the relationship between presence of *M. lehtineni* and the specific microclimate caused by the melting snow in the mountains was also suggested (Bolzern et al. 2005).

The biology of the presented species is poorly known, because it is extremely rare and difficult to observe. Hence, any new data on its distribution are of considerable importance.



**Fig. 6-8:** *Maro lehtineni*, lateral view of palpal organs of the two males. **6-7.** first male, **8.** second male. The shape of the lamella characteristic is changeable and depends on the viewing angle. Arrow points to the lamella characteristic.

## References

- Baert LL & Kekenbosch J 1979 Le genre *Maro* O.P.-Cambridge 1906 (Araneae: Linyphiidae) en Belgique. – Bulletin et Annales de la Société Royale Belge d'Entomologie 115: 117-120
- Blick T, Bosmans R, Buchar J, Gajdoš P, Hänggi A, Helsdingen P van, Růžicka V, Starega W & Thaler K 2004 Checkliste der Spinnen Mitteleuropas. Checklist of the spiders of Central Europe. (Arachnida: Araneae). Version 1. December 2004. – Internet: [http://arages.de/wp-content/uploads/2013/05/checklist2004\\_araneae.pdf](http://arages.de/wp-content/uploads/2013/05/checklist2004_araneae.pdf) (June 5, 2015)
- Bolzern A, Hänggi A, Kropf C & Frick H 2005 Erstnachweis von *Maro lehtineni* Saaristo, 1971 für die Schweiz (Arachnida, Araneae). – Mitteilungen der Schweizerischen entomologischen Gesellschaft 78: 165-172
- Casimir H 1976 Beitrag zur Hochmoor-Spinnenfauna des Hohen Venns (Hautes Fagnes) zwischen Nordeifel und Ardennen. – Decheniana 129: 38-72
- Granström U 1978 The spider fauna of the province Västernorrland. A species list with faunistic and phenological notes. – Fauna Norrlandica 10: 1-14
- Hauge E 1980 A contribution to the spider fauna (Araneae) of Norway. – Fauna norvegica B 27: 68-71
- Helsdingen P van 2015 Fauna Europaea: Spiders. Fauna Europaea version 2.6.2. – Internet: <http://www.faunaeur.org> (June 5, 2015)
- Kupryjanowicz J 2003 Spiders (Araneae) of open habitats in the Biebrza National Park, Poland. – Fragmenta Faunistica, Warsaw 46: 209-237
- Kupryjanowicz J 2008 Pająki, Araneae. In: Bogdanowicz W, Chudzicka E, Pilipiuk I & Skibińska E (eds) Fauna Polski. Charakterystyka i wykaz gatunków. Muzeum i Instytut Zoologii PAN, Warszawa. pp. 223-239
- Martin D 2009 *Ceraticelus bulbosus* (Araneae, Linyphiidae) – Erstnachweis für Deutschland sowie weitere bemerkenswerte Spinnenfunde aus Ostdeutschland. – Arachnologische Mitteilungen 38: 4-7 – doi: 10.5431/aramit3802
- Muff P, Schmidt MH, Frick H, Nentwig W & Kropf C 2007 Spider (Arachnida: Araneae) distribution across the timberline in the Swiss Central Alps (Alp Flix, Grisons) and three morphologically remarkable species. – Arachnologische Mitteilungen 34: 16-24 – doi: 10.5431/aramit3404
- Palmgren P 1975 Die Spinnenfauna Finnland und Ostfennoskandiens VI. Linyphiidae 1 (Die Linyphiinae und Linyphiinae-ähnlichen Micryphantinae). – Fauna Fennica 28: 1-102
- Růžicka V 2000 Spiders (Araneae) of two valleys in the Krkonoše Mts. (Czech Republic). In: Pekár S (ed.) Proceedings of the 18th European Colloquium of Arachnology, 1999. – Ekologia (Bratislava) 19, Supplement 3: 235-244
- Saaristo MI 1971 Revision of the genus *Maro* O. P.-Cambridge (Araneae, Linyphiidae). – Annales Zoologici Fennici 8: 463-482
- Stańska M, Hajdamowicz I & Żabka M 2002 Epigeic spiders of alder swamp forests in Eastern Poland. In: Toft S & Scharff N (ed.) European Arachnology 2000. Aarhus University Press, Aarhus. pp. 191-197
- Starega W 1996 Spinnen (Araneae) aus der Borkenheide und anderen Lokalitäten der Masurischen Seenplatte. – Fragmenta faunistica, Warsaw 39: 287-311
- Starega W 2000 Spinnen (Araneae) aus Roztocze und den anliegenden Gebieten. – Fragmenta Faunistica, Warsaw 47: 59-89
- Svatoň J & Gajdoš P 2004 Spiders of peatland ecosystems of the Horná Orava region (Slovakia). In: Szinetár C (ed.) European Arachnology 2002. Plant Protection Institute & Berzsenyi College, Budapest. pp. 275-284
- Szymkowiak P 2004 *Maro sublestus* Falconer, 1915 (Araneae, Linyphiidae) – a spider new to the fauna of Poland. – Fragmenta Faunistica, Warsaw 47: 139-142
- Tanasevitch AV & Nekhaeva AA 2014 Predvaritel'nye itogi araneologicheskikh issledovaniy v Pinezhskom zapovednike (Arachnida, Aranei). [Preliminary results of araneological studies in the Pinezhsky Nature Reserve (Arachnida, Aranei)]. In: Putschina LV (ed.), Sokhranenie i izutchenie geo- i bioraznობrazii na OOPT Evropeiskogo Severa Rossii. Materialy nauchno-prakticheskoi konferencii, posviashchennoi 40-letiiu zapovednika „Pinezhskii”, 2-5.IX.2014. Pinega, Arkhangiel'skaia oblast', Izhevsk. pp. 204-207 [in Russian]
- Thaler K 1983 Bemerkenswerte Spinnenfunde in Nordtirol (Österreich) und Nachbarländern: Deckennetzspinnen, Linyphiidae (Arachnida, Aranei). – Veröffentlichungen des Museum Ferdinandeum in Innsbruck 63: 135-167
- Wiśniewski K & Wesołowska W 2012 *Maro lepidus* Casimir, 1961, a newly recorded spider species (Araneae, Linyphiidae) for Poland. – Fragmenta Faunistica, Warsaw 55: 155-160
- World Spider Catalog 2015 World spider catalog, version 16. Natural History Museum, Bern. – Internet: <http://wsc.nmbe.ch> (June 5, 2015)
- Woźny M, Czajka M, Pilawski S & Bednarz S 1988 Pająki (Aranei) polskich Sudetów. – Acta Universitatis Wratislaviensis 972, Prace Zoologiczne 19: 53-130